

# 3-Abelian Cubes Are Avoidable on Binary Alphabets

Robert Mercas<sup>1\*</sup> and Aleksi Saarela<sup>2\*\*</sup>

<sup>1</sup> Otto-von-Guericke-Universität Magdeburg, Fakultät für Informatik,  
PSF 4120, D-39016 Magdeburg, Germany,

`robertmercas@gmail.com`

<sup>2</sup> Department of Mathematics and Statistics, University of Turku,  
FI-20014 Turku, Finland,

`amsaar@utu.fi`

**Abstract.** A  $k$ -abelian cube is a word  $uvw$ , where  $u, v, w$  have the same factors of length at most  $k$  with the same multiplicities. Previously it has been known that  $k$ -abelian cubes are avoidable over a binary alphabet for  $k \geq 5$ . Here it is proved that this holds for  $k \geq 3$ .

**Keywords:** combinatorics on words, repetition-freeness,  $k$ -abelian equivalence

## 1 Introduction

The study of repetition-free infinite words (or even the whole area of combinatorics on words) was begun by Axel Thue [1, 2]. He proved that using three letters one can construct an infinite word that does not contain a square, that is a factor of the form  $uu$  where  $u$  is a non-empty word. Further, using two letters one can construct an infinite word that does not contain a cube, that is a factor of the form  $uuu$  where  $u$  is a non-empty word, or even an overlap, that is a factor of the form  $auaua$  where  $u$  is a word and  $a$  is a letter. Due to their initial obscure publication, these results have been rediscovered several times.

The problem of repetition-freeness has been studied from many points of view. One is to consider fractional powers. This leads to the concept of repetition threshold and the famous Dejean's conjecture, which was proved in many parts, of which the last one independently in [3] and [4]. Another example is the repetition-freeness of partial words. It was shown that there exists infinite ternary words with an infinite number of holes whose factors are not matching any squares (overlaps) of words of length greater than one [5, 6]. For the abelian case an alphabet with as low as 5 letters is enough in order to construct an infinite word with factors that do not match an abelian square of any word of length greater than two [7].

---

\* Supported by the Alexander von Humboldt Foundation

\*\* Supported by the Academy of Finland under grant 257857

In this paper abelian repetition-freeness is an important concept. Abelian square is a non-empty word  $uv$ , where  $u$  and  $v$  have the same number of occurrences of each symbol. Abelian cubes and  $n$ th powers are defined in a similar way. Erdős [8] raised the question whether abelian squares can be avoided, i.e., whether there exist infinite words over a given alphabet that do not contain two consecutive permutations of the same factor. It is easily seen that abelian squares cannot be avoided over a three-letter alphabet: Each word of length eight over three letters contains an abelian square. Dekking [9] proved that over a binary alphabet there exists a word that avoids abelian fourth powers, and over a ternary alphabet there exists a word that avoids abelian cubes. The problem of whether abelian squares can be avoided over a four-letter alphabet was open for a long time. In [10], using an interesting combination of computer checking and mathematical reasoning, Keränen proved that abelian squares are avoidable on four letters.

Recently, several questions have been studied from the point of view of  $k$ -abelian equivalence. For a positive integer  $k$ , two words are said to be  $k$ -abelian equivalent if they have the same number of occurrences of every factor of length at most  $k$ . These equivalence relations provide a bridge between abelian equivalence and equality, because 1-abelian equivalence is the same as abelian equivalence, and as  $k$  grows,  $k$ -abelian equivalence becomes more and more like equality. The topic of this paper is  $k$ -abelian repetition-freeness, but there has also been research on other topics related to  $k$ -abelian equivalence [11, 12].

In [11], the authors show that 2-abelian squares are avoidable only on a four letter alphabet. For  $k \geq 3$ , the question of avoiding  $k$ -abelian squares is open, the minimal alphabet size being either three or four. Computer experiments would suggest that there are 3-abelian square-free ternary words, but it is known that there are no pure morphic  $k$ -abelian square-free ternary words for any  $k$  [13].

It was conjectured in [11] that for avoiding  $k$ -abelian cubes a binary alphabet suffices whenever  $k \geq 2$  since computer generated words of length 100000 having the property have been found. This was proved for  $k \geq 8$  in [14] and for  $k \geq 5$  in [15].

In this work it is proved that 3-abelian cubes can be avoided on a binary alphabet. The methods used are somewhat similar to those used in [14] and [15]: A word is constructed by mapping an abelian cube-free ternary word by a morphism. However, there are some crucial differences. Most importantly, the morphisms used in this paper are not uniform, and this makes many parts of the proofs different and more difficult. The method used in this article is fairly general, but using it requires an extensive case analysis, which can only be carried out with the help of a computer. The 2-abelian case remains open.

## 2 Preliminaries

We denote by  $\Sigma$  a finite set of symbols called *alphabet*. For  $n \geq 0$ , the  $n$ -letter alphabet  $\{0, \dots, n-1\}$  will be denoted by  $\Sigma_n$ . A *word*  $w$  represents a concatenation of letters from  $\Sigma$ . By  $\varepsilon$  we denote the *empty word*. We denote

by  $|w|$  the *length* of  $w$  and by  $|w|_u$  the number of occurrences of  $u$  in  $w$ . The *concatenation* of two words  $u$  and  $v$  is the word  $uv$  obtained by adding to the right of  $u$  the letters of the word  $v$ . For a factorization  $w = uxv$ , we say that  $x$  is a *factor* of  $w$ , and whenever  $u$  is empty  $x$  is a *prefix* and, respectively, when  $v$  is empty  $x$  is a *suffix* of  $w$ .

We will use the following notation for factors, prefixes and suffixes. If  $w = a_0a_1 \dots a_{n-1}$  with  $a_i \in \Sigma$  for  $0 \leq i < n$  we say that  $\text{rfact}_k^i(w) = a_i \dots a_{i+k-1}$  is the *right factor at  $i$*  of  $w$  of length  $k$  and that  $\text{lfact}_k^i(w) = a_{i-k} \dots a_{i-1}$  is the *left factor at  $i$*  of  $w$  of length  $k$ . Furthermore,  $\text{pref}_k(w) = \text{rfact}_k^0(w)$  is the length  $k$  *prefix* of  $w$  and  $\text{suff}_k(w) = \text{lfact}_k^n(w)$  is the length  $k$  *suffix* of  $w$ .

The *powers* of a word  $w$  are defined recursively,  $w^0 = \varepsilon$  and  $w^n = ww^{n-1}$  for  $n > 0$ . We say that  $w$  is an  *$n$ th power* if there exists a word  $u$  such that  $w = u^n$ . Second powers are called *squares* and third powers *cubes*.

Words  $u$  and  $v$  are *abelian equivalent* if  $|u|_a = |v|_a$  for all letters  $a \in \Sigma$ . For a word  $u \in \Sigma_n^*$ , let  $P_u = (|u|_0, \dots, |u|_{n-1})$  be the *Parikh vector* of  $u$ . Words  $u, v \in \Sigma_n^*$  are abelian equivalent if and only if  $P_u = P_v$ .

Two words  $u$  and  $v$  are  *$k$ -abelian equivalent* if  $|u|_t = |v|_t$  for every word  $t$  of length at most  $k$ . Obviously, 1-abelian equivalence is the same as abelian equivalence, and words of length less than  $k - 1$  (or, in fact, words of length less than  $2k$ ) are equivalent only if they are equal. For words  $u$  and  $v$  of length at least  $k - 1$ , another equivalent definition can be given:  $u$  and  $v$  are  *$k$ -abelian equivalent* if  $|u|_t = |v|_t$  for every word  $t$  of length  $k$ ,  $\text{pref}_{k-1}(u) = \text{pref}_{k-1}(v)$  and  $\text{suff}_{k-1}(u) = \text{suff}_{k-1}(v)$ . This latter definition is actually the one used in the proofs of this article.

A  *$k$ -abelian  $n$ th power* is a word  $u_1u_2 \dots u_n$ , where  $u_1, u_2, \dots, u_n$  are  *$k$ -abelian equivalent*. For  $k = 1$  this gives the definition of an *abelian  $n$ th power*.

A mapping  $f : A^* \rightarrow B^*$  is a *morphism* if  $f(xy) = f(x)f(y)$  for any words  $x, y \in A^*$ , and is completely determined by the images  $f(a)$  for all  $a \in A$ .

If no non-empty square is a factor of a word  $w$ , then it is said that  $w$  is *square-free*, or that  $w$  *avoids squares*. If there exists an infinite square-free word over an alphabet  $\Sigma$ , then it is said that *squares are avoidable on  $\Sigma$* . Of course the only thing that matters here is the size of  $\Sigma$ . Similar definitions can be given for cubes and higher powers, as well as for  *$k$ -abelian powers*.

Unlike ordinary cubes, abelian cubes are nor avoidable on a binary alphabet, and unlike ordinary squares, abelian squares are not avoidable on a ternary alphabet. However, Dekking showed in [9] that two letters are sufficient for avoiding abelian fourth powers, and three letters suffice for avoiding abelian cubes. An extension of the latter result is stated in the following theorem. It is proved that the word of Dekking avoids also many other factors in addition to abelian cubes.

**Theorem 1.** *Let  $w$  be the fixed point of the morphism  $\sigma : \Sigma_3^* \rightarrow \Sigma_3^*$  defined by*

$$\sigma(0) = 0012, \quad \sigma(1) = 112, \quad \sigma(2) = 022.$$

*Then  $w$  is abelian cube-free and contains no factor  $apbqbrc$  where one of the following conditions is satisfied:*

1.  $abcd = 0112$  and  $P_p = P_q = P_r$ ,
2.  $abcd = 0210$  and  $P_p = P_q - (1, -1, 1) = P_r - (0, -1, 1)$ ,
3.  $abcd = 0211$  and  $P_p = P_q - (1, -1, 1) = P_r - (1, -2, 1)$ ,
4.  $abcd = 0220$  and  $P_p = P_q - (1, -1, 1) = P_r - (0, 0, 0)$ ,
5.  $abcd = 0221$  and  $P_p = P_q - (1, -1, 1) = P_r - (1, -1, 0)$ ,
6.  $abcd = 1001$  and  $P_p = P_q = P_r$ ,
7.  $abcd = 1002$  and  $P_p = P_q = P_r$ ,

*Proof.* The word  $w$  was shown to be abelian cube-free in [9]. Similar ideas can be used to show that  $w$  avoids the factors  $apbqcrd$ . Case 1 was proved in [15]. Case 2 is proved here. Cases 3–6 are similar as the first two, so their proofs are omitted. Case 7 is more difficult, so it is proved here.

Let  $f : \Sigma^* \rightarrow \mathbb{Z}_7$  be the morphism defined by

$$f(0) = 1, \quad f(1) = 2, \quad f(2) = 3$$

(here  $\mathbb{Z}_7$  is the additive group of integers modulo 7). Then  $f(\sigma(x)) = 0$  for all  $x \in \Sigma$ . If  $apbqcrd$  is a factor of  $w$ , then there are  $u, s, t$  such that  $\sigma(u) = sapbqcrdt$  and  $u$  is a factor of  $w$ . Consider the values

$$f(s), f(sa), f(sap), f(sapb), f(sapbq), f(sapbqc), f(sapbqcr), f(sapbqcrd). \quad (1)$$

These elements are of the form  $f(\sigma(u')v') = f(v')$ , where  $v'$  is a prefix of one of 0012, 112, 022. The possible values for  $f(v')$  are 0, 1, 2 and 4.

Consider Case 2. Let  $abcd = 0210$ . If  $P_p = P_q - (1, -1, 1) = P_r - (0, -1, 1)$ , then  $f(p) = f(q) - 2 = f(r) - 1$ . If we denote  $i = f(s)$ ,  $j = f(p)$ , then the values for (1) are

$$i, i + 1, i + j + 1, i + j + 4, i + 2j + 6, i + 2j + 1, i + 3j + 2, i + 3j + 3.$$

For all values of  $i$  and  $j$ , one is not 0, 1, 2 or 4. This is a contradiction.

Consider Case 7. Let  $abcd = 1002$ . If  $P_p = P_q = P_r$ , then  $f(p) = f(q) = f(r)$ . If we denote  $i = f(s)$ ,  $j = f(p)$ , then the values for (1) are

$$i, i + 2, i + j + 2, i + j + 3, i + 2j + 3, i + 2j + 4, i + 3j + 4, i + 3j.$$

It must be  $i = 0$  and  $j = 6$ , because otherwise one of the values is not 0, 1, 2 or 4. There are letters  $a', b', c', d'$  and words  $s', p', q', r', t', s_2, p_1, p_2, q_1, q_2, r_1, r_2, t_1$  such that

$$\begin{array}{ll} u = s'a'p'b'q'r'd't' & s = \sigma(s')s_2 \\ s_21p_1 = \sigma(a') & p = p_1\sigma(p')p_2 \\ p_20q_1 = \sigma(b') & q = q_1\sigma(q')q_2 \\ q_20r_1 = \sigma(c') & r = r_1\sigma(r')r_2 \\ r_22t_1 = \sigma(d') & t = t_1\sigma(t'), \end{array}$$

i.e. the situation is like in the following diagram:

|              |              |              |              |              |              |              |              |              |
|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| $s$          | $ 1 $        | $p$          | $ 0 $        | $q$          | $ 0 $        | $r$          | $ 2 $        | $t$          |
|              | $s_2$        | $p_1$        | $p_2$        | $q_1$        | $q_2$        | $r_1$        | $r_2$        | $t_1$        |
| $\sigma(s')$ | $\sigma(a')$ | $\sigma(p')$ | $\sigma(b')$ | $\sigma(q')$ | $\sigma(c')$ | $\sigma(r')$ | $\sigma(d')$ | $\sigma(t')$ |

Because  $i = 0$ ,  $s_2 = \varepsilon$ . Then  $\sigma(a')$  begins with 1, so  $a' = 1$  and  $p_1 = 12$ . Thus  $p = 12\sigma(p')p_2$ . It must be  $f(p_2) = f(p) - f(\sigma(p')) - f(12) = j - 0 - 5 = 1$ , so  $p_2 = 0$ . Then  $\sigma(b')$  begins with 00, so  $b' = 0$  and  $q_1 = 12$ . Like above, it can be concluded that  $q = 12\sigma(q')0$ , and similarly also  $r = 12\sigma(r')0$ . But then  $1p'0q'0r'2$  is a factor of  $w$ . If

$$M = \begin{pmatrix} |\sigma(0)|_0 & |\sigma(1)|_0 & |\sigma(2)|_0 \\ |\sigma(0)|_1 & |\sigma(1)|_1 & |\sigma(2)|_1 \\ |\sigma(0)|_2 & |\sigma(1)|_2 & |\sigma(2)|_2 \end{pmatrix} = \begin{pmatrix} 2 & 1 & 1 \\ 0 & 2 & 1 \\ 1 & 0 & 2 \end{pmatrix}$$

and Parikh vectors are interpreted as column vectors, then

$$MP_{p'} = P_{\sigma(p')}, \quad MP_{q'} = P_{\sigma(q')}, \quad MP_{r'} = P_{\sigma(r')}.$$

Because  $M$  is invertible and  $\sigma(p'), \sigma(q'), \sigma(r')$  are abelian equivalent, also  $p', q', r'$  are abelian equivalent. By repeating the process, we get shorter and shorter factors of the same form, which leads to a contradiction.  $\square$

If abelian cubes are avoidable on some alphabet, then so are  $k$ -abelian cubes. This means that  $k$ -abelian cubes are avoidable on a ternary alphabet for all  $k$ . But for which  $k$  are they avoidable on a binary alphabet? In [14] it was proved that this holds for  $k \geq 8$ , and conjectured that it holds for  $k \geq 2$ . In [15] it was proved that this holds for  $k \geq 5$ . In this article it is proved that this holds for  $k \geq 3$ . The case when  $k = 2$  remains open.

### 3 3-abelian cube-freeness

Let  $w \in \Sigma_m^\omega$ . The following remarks will be used in the case where  $m = 3$ ,  $n = 2$ ,  $w$  is abelian cube-free and  $k = 4$  or  $k = 3$ , but they hold also more generally.

For a word  $v \in \Sigma_n^*$ , let  $Q_v = (|v|_{t_0}, \dots, |v|_{t_{N-1}})$ , where  $t_0, \dots, t_{N-1}$  are the words of  $\Sigma_n^k$  in lexicographic order. When doing matrix calculations, all vectors  $P_u$  and  $Q_v$  will be interpreted as column vectors.

Let  $h : \Sigma_m^* \rightarrow \Sigma_n^*$  be a morphism. It needs to be assumed that  $h$  satisfies three conditions:

- There is a word  $s \in \Sigma_m^{k-1}$  that is a prefix of  $h(a)$  for every  $a \in \Sigma_m$ .
- The matrix  $M$  whose columns are  $Q_{h(0)s}, \dots, Q_{h(m-1)s}$  has rank  $m$ .
- There are no  $k$ -abelian equivalent words  $v_1, v_2, v_3$  of length less than

$$2 \max \{h(a) \mid a \in \Sigma_m\}$$

such that  $v_1 v_2 v_3$  is a factor of  $h(w)$ .

Let  $M^+$  be the Moore-Penrose pseudoinverse of  $M$ . The only properties of  $M^+$  needed in this article are that it exists and can be efficiently computed, and that since the columns of  $M$  are linearly independent,  $M^+M$  is the  $m \times m$  identity matrix. For any word  $u \in \Sigma^*$ ,  $Q_{h(u)s} = MP_u$ .

**Lemma 2.** *If the word  $h(w)$  has a factor  $v_1v_2v_3$ , where  $v_1, v_2, v_3$  are  $k$ -abelian equivalent, then there are letters  $a_0, a_1, a_2, a_3, b_2, b_3 \in \Sigma_m$ , words  $u_1, u_2, u_3 \in \Sigma_m^*$  and indices*

$$\begin{aligned} i_0 &\in \{0, \dots, |h(a_0)| - 1\}, \\ i_1 &\in \{k - 1, \dots, |h(a_1)| + k - 2\}, \\ i_2 &\in \{k - 1, \dots, |h(a_2)| + k - 2\}, \\ i_3 &\in \{k - 1, \dots, |h(a_3)| + k - 2\} \end{aligned} \tag{2}$$

such that  $a_0u_1a_1b_2u_2a_2b_3u_3a_3$  is a factor of  $w$  and  $v_i = x_ih(u_i)y_i$  for  $i \in \{1, 2, 3\}$ , where

$$\begin{aligned} x_1 &= \text{suff}_{|h(a_0)|-i_0}(h(a_0)) & y_1 &= \text{pref}_{i_1}(h(a_1b_2)), \\ x_2 &= \text{suff}_{|h(a_1b_2)|-i_1}(h(a_1b_2)) & y_2 &= \text{pref}_{i_2}(h(a_2b_3)), \\ x_3 &= \text{suff}_{|h(a_2b_3)|-i_2}(h(a_2b_3)) & y_3 &= \text{pref}_{i_3}(h(a_3)s). \end{aligned} \tag{3}$$

*Proof.* It was assumed that  $h(w)$  does not contain short  $k$ -abelian cubes, and a longer  $k$ -abelian cube  $v_1v_2v_3$  must be of the form specified in the claim.  $\square$

Because  $s$  is a prefix of  $h(u_i)$  and  $y_i$ , it follows that  $Q_{v_i} = Q_{x_i s} + Q_{h(u_i)s} + Q_{y_i}$ .

The idea is to iterate over all values of  $a_0, a_1, a_2, a_3, b_2, b_3$  and  $i_0, i_1, i_2, i_3$  and in each case try to deduce that one of the following holds:

- There are no  $u_1, u_2, u_3$  such that the words  $v_i = x_ih(u_i)y_i$  are  $k$ -abelian equivalent.
- If  $v_i = x_ih(u_i)y_i$  are  $k$ -abelian equivalent, then  $a_0u_1a_1b_2u_2a_2b_3u_3a_3$  contains an abelian cube or a factor of the form mentioned in Theorem 1.

If we succeed, then there are words  $w$  such that  $h(w)$  is  $k$ -abelian cube-free. The following lemmas will be useful.

**Lemma 3.** *Let  $a_0, a_1, a_2, a_3, b_2, b_3 \in \Sigma_m$ , indices  $i_0, i_1, i_2, i_3$  be as in (2) and words  $x_1, x_2, x_3, y_1, y_2, y_3$  be as in (3). If*

$$\begin{aligned} \text{pref}_{k-1}(x_1), \text{pref}_{k-1}(x_2), \text{pref}_{k-1}(x_3) &\text{ are not equal or} \\ \text{suff}_{k-1}(y_1), \text{suff}_{k-1}(y_2), \text{suff}_{k-1}(y_3) &\text{ are not equal,} \end{aligned} \tag{C1}$$

then there are no  $u_1, u_2, u_3$  such that the three words  $v_i = x_ih(u_i)y_i$  would be  $k$ -abelian equivalent.

*Proof.* If the prefixes or suffixes of  $v_1, v_2, v_3$  of length  $k - 1$  are not equal, then  $v_1, v_2, v_3$  cannot be  $k$ -abelian equivalent.  $\square$

**Lemma 4.** Let  $a_0, a_1, a_2, a_3, b_2, b_3 \in \Sigma_m$ , indices  $i_0, i_1, i_2, i_3$  be as in (2) and words  $x_1, x_2, x_3, y_1, y_2, y_3$  be as in (3). Let  $R_i = Q_{x_{i_s}} + Q_{y_i}$  for  $i \in \{1, 2, 3\}$ . If

$$\begin{aligned} M^+(R_1 - R_i) \text{ is not an integer vector or} \\ MM^+(R_1 - R_i) + R_i \text{ are not equal for } i \in \{1, 2, 3\}, \end{aligned} \quad (\text{C2})$$

then there are no  $u_1, u_2, u_3$  such that the three words  $v_i = x_i h(u_i) y_i$  would be  $k$ -abelian equivalent.

*Proof.* If  $v_i = x_i h(u_i) y_i$ , then  $Q_{v_i} = Q_{h(u_i)s} + R_i = MP_{u_i} + R_i$ . If  $Q_{v_1} = Q_{v_2} = Q_{v_3}$ , then  $P_{u_i} - P_{u_1} = M^+(R_1 - R_i)$ . This must be an integer vector. The vectors  $Q_{v_i} - MP_{u_1} = MM^+(R_1 - R_i) + R_i$  must be equal for  $i \in \{1, 2, 3\}$ .  $\square$

**Lemma 5.** Let  $a_0, a_1, a_2, a_3, b_2, b_3 \in \Sigma_m$ , indices  $i_0, i_1, i_2, i_3$  be as in (2) and words  $x_1, x_2, x_3, y_1, y_2, y_3$  be as in (3). Let  $R_i = Q_{x_{i_s}} + Q_{y_i}$  for  $i \in \{1, 2, 3\}$ . If

$$\begin{aligned} \text{for } i \in \{0, 1, 2, 3\} \text{ there are } c_i, d_i \in \{a_i, \varepsilon\} \text{ such that } c_i d_i = a_i \text{ and} \\ M^+(R_1 - R_1) + P_{d_0 c_1} \\ = M^+(R_1 - R_2) + P_{d_1 b_2 c_2} \\ = M^+(R_1 - R_3) + P_{d_2 b_3 c_3} \end{aligned} \quad (\text{C3})$$

and  $a_0 u_1 a_1 b_2 u_2 a_2 b_3 u_3 a_3$  is abelian cube-free, then the three words  $v_i = x_i h(u_i) y_i$  cannot be  $k$ -abelian equivalent.

*Proof.* Like in the proof of Lemma 4, the  $k$ -abelian equivalence of  $v_1, v_2, v_3$  implies  $P_{u_i} - P_{u_1} = M^+(R_1 - R_i)$ . From this and (C3) it follows that

$$P_{u_1} + P_{d_0 c_1} = P_{u_2} + P_{d_1 b_2 c_2} = P_{u_3} + P_{d_2 b_3 c_3},$$

so  $d_0 u_1 c_1, d_1 b_2 u_2 c_2, d_2 b_3 u_3 c_3$  are abelian equivalent. This contradicts the abelian cube-freeness of  $a_0 u_1 a_1 b_2 u_2 a_2 b_3 u_3 a_3$ .  $\square$

**Lemma 6.** Let  $a_0, a_1, a_2, a_3, b_2, b_3 \in \Sigma_m$ , indices  $i_0, i_1, i_2, i_3$  be as in (2) and words  $x_1, x_2, x_3, y_1, y_2, y_3$  be as in (3). Let  $R_i = Q_{x_{i_s}} + Q_{y_i}$  for  $i \in \{1, 2, 3\}$  and  $S_i = M^+(R_1 - R_i) + P_{b_i}$  for  $i \in \{2, 3\}$ . If

$$\begin{aligned} (0 = S_2 = S_3) & \quad \text{and} \quad a_0 a_1 a_2 a_3 = 0112) \text{ or} \\ (0 = S_2 - (1, -1, 1) = S_3 - (0, -1, 1)) & \quad \text{and} \quad a_0 a_1 a_2 a_3 = 0210) \text{ or} \\ (0 = S_2 - (1, -1, 1) = S_3 - (1, -2, 1)) & \quad \text{and} \quad a_0 a_1 a_2 a_3 = 0211) \text{ or} \\ (0 = S_2 - (1, -1, 1) = S_3 - (0, 0, 0)) & \quad \text{and} \quad a_0 a_1 a_2 a_3 = 0220) \text{ or} \\ (0 = S_2 - (1, -1, 1) = S_3 - (1, -1, 0)) & \quad \text{and} \quad a_0 a_1 a_2 a_3 = 0221) \text{ or} \\ (0 = S_2 = S_3) & \quad \text{and} \quad a_0 a_1 a_2 a_3 = 1001) \text{ or} \\ (0 = S_2 = S_3) & \quad \text{and} \quad a_0 a_1 a_2 a_3 = 1002) \end{aligned} \quad (\text{C4})$$

and  $a_0 u_1 a_1 b_2 u_2 a_2 b_3 u_3 a_3$  is not of the form  $apbqcrd$  specified in Theorem 1, then the three words  $v_i = x_i h(u_i) y_i$  cannot be  $k$ -abelian equivalent.

*Proof.* Like in the proof of Lemma 4, the  $k$ -abelian equivalence of  $v_1, v_2, v_3$  implies  $P_{u_i} - P_{u_1} = M^+(R_1 - R_i)$ . From this and the first row of (C4) it follows that

$$P_{u_1} = P_{u_2} + P_{b_2} = P_{u_3} + P_{b_3},$$

so  $u_1, b_2u_2, b_3u_3$  are abelian equivalent, which is a contradiction. The other rows lead to a contradiction in a similar way.  $\square$

We can iterate over all values of  $a_0, a_1, a_2, a_3, b_2, b_3$  and  $i_0, i_1, i_2, i_3$ . If in all cases one of Conditions C1, C2, C3 is true, then  $h$  maps all abelian cube-free words to  $k$ -abelian cube-free words. If in all cases one of Conditions C1, C2, C3, C4 is true, then  $h$  maps the word of Theorem 1 to a  $k$ -abelian cube-free word. In this way we obtain Theorems 7 and 8.

Concerning the actual implementation of the above algorithm, there are some optimizations that can be made. First, if  $i_1$  and  $i_2$  are such that  $b_1$  and  $b_2$  do not affect the definition of  $x_1, x_2, x_3, y_1, y_2, y_3$  in (3), then instead of iterating over all values of  $b_1$  and  $b_2$ , they can be combined with  $u_2$  and  $u_3$ . Second, in most of the cases Condition C1 is true, and these cases can be handled easily. In the following theorems, there are a couple of thousand nontrivial cases, i.e. cases where Condition C1 is false.

**Theorem 7.** *The morphism defined by*

$$0 \mapsto 10110100110, \quad 1 \mapsto 101101001001, \quad 2 \mapsto 1011001100100,$$

*maps every abelian cube-free ternary word to a 4-abelian cube-free word.*

**Theorem 8.** *The morphism defined by*

$$0 \mapsto 01010, \quad 1 \mapsto 0110010, \quad 2 \mapsto 0110110,$$

*maps the word  $w$  of Theorem 1 to a 3-abelian cube-free word.*

## References

1. Thue, A.: Über unendliche Zeichenreihen. Norske Vid. Selsk. Skr. I, Mat. Nat. Kl. Christiania **7** (1906) 1–22 (Reprinted in *Selected Mathematical Papers of Axel Thue*, T. Nagell, editor, Universitetsforlaget, Oslo, Norway (1977), pp. 139–158).
2. Thue, A.: Über die gegenseitige Lage gleicher Teile gewisser Zeichenreihen. Norske Vid. Selsk. Skr. I, Mat. Nat. Kl. Christiania **1** (1912) 1–67 (Reprinted in *Selected Mathematical Papers of Axel Thue*, T. Nagell, editor, Universitetsforlaget, Oslo, Norway (1977), pp. 413–478).
3. Currie, J., Rampersad, N.: A proof of Dejean’s conjecture. *Math. Comp.* **80** (2011) 1063–1070
4. Rao, M.: Last cases of Dejean’s conjecture. *Theoretical Computer Science* **412**(27) (2011) 3010–3018
5. Manea, F., Mercas, R.: Freeness of partial words. *Theoretical Computer Science* **389**(1-2) (2007) 265–277



6. Blanchet-Sadri, F., Mercas, R., Scott, G.: A generalization of Thue freeness for partial words. *Theoretical Computer Science* **410**(8-10) (2009) 793–800
7. Blanchet-Sadri, F., Kim, J.I., Mercas, R., Severa, W., Simmons, S., Xu, D.: Avoiding abelian squares in partial words. *Journal of Combinatorial Theory, Series A* **119**(1) (2012) 257 – 270
8. Erdős, P.: Some unsolved problems. *Magyar Tudományos Akadémia Matematikai Kutató Intézete* **6** (1961) 221–254
9. Dekking, F.M.: Strongly nonrepetitive sequences and progression-free sets. *J. Combin. Theory Ser. A* **27**(2) (1979) 181–185
10. Keränen, V.: Abelian squares are avoidable on 4 letters. In: *Proceedings of the 19th International Colloquium on Automata, Languages and Programming.* (1992) 41–52
11. Huova, M., Karhumäki, J., Saarela, A., Saari, K.: Local squares, periodicity and finite automata. In Calude, C., Rozenberg, G., Salomaa, A., eds.: *Rainbow of Computer Science.* Springer (2011) 90–101
12. Karhumäki, J., Puzynina, S., Saarela, A.: Fine and Wilf’s theorem for  $k$ -abelian periods. In: *Proceedings of the 16th International Conference on Developments in Language Theory.* (2012) 296–307
13. Huova, M.: On unavailability of  $k$ -abelian squares in pure morphic words. In: *Proceedings of the 14th Mons Days of Theoretical Computer Science.* (2012)
14. Huova, M., Karhumäki, J., Saarela, A.: Problems in between words and abelian words:  $k$ -abelian avoidability. *Theoret. Comput. Sci.* **454** (2012) 172–177
15. Mercas, R., Saarela, A.: 5-abelian cubes are avoidable on binary alphabets. In: *Proceedings of the 14th Mons Days of Theoretical Computer Science.* (2012)